



Draft genome sequence of *Chelonobacter oris* strain 1662T, associated with respiratory disease in Hermann's tortoises

Kudirkiene, Egle; Hansen, Mie Johanne; Bojesen, Anders Miki

Published in:
Genome Announcements

DOI:
[10.1128/genomeA.01322-14](https://doi.org/10.1128/genomeA.01322-14)

Publication date:
2014

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Kudirkiene, E., Hansen, M. J., & Bojesen, A. M. (2014). Draft genome sequence of *Chelonobacter oris* strain 1662^T, associated with respiratory disease in Hermann's tortoises. *Genome Announcements*, 2(6), [e01322-14]. <https://doi.org/10.1128/genomeA.01322-14>

Draft Genome Sequence of *Chelonobacter oris* Strain 1662^T, Associated with Respiratory Disease in Hermann's Tortoises

Egle Kudirkienė, Mie Johanne Hansen, Anders Miki Bojesen

Department of Veterinary Disease Biology, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

***Chelonobacter oris* 1662^T is a type strain of the recently described species of the *Pasteurellaceae* family. The strain was isolated from the choanae of a captive tortoise with signs of respiratory tract infection. The genome reported here is approximately 2.6 Mb in size and has a G+C content of 47.1%.**

Received 10 November 2014 Accepted 11 November 2014 Published 18 December 2014

Citation Kudirkienė E, Hansen MJ, Bojesen AM. 2014. Draft genome sequence of *Chelonobacter oris* strain 1662^T, associated with respiratory disease in Hermann's tortoises. *Genome Announc.* 2(6):e01322-14. doi:10.1128/genomeA.01322-14.

Copyright © 2014 Kudirkienė et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Anders Miki Bojesen, miki@sund.ku.dk.

Pasteurellaceae, with few exceptions, are primarily opportunistic pathogens. Until 2011, the *Pasteurellaceae* family included 15 genera (1); however, new genera isolated from various animal species are continuously added to the family *Pasteurellaceae* (2–6). Most *Pasteurellaceae* species are host-specific and only occasionally may be transmitted between different hosts (7).

Chelonobacter oris, currently the only species in the *Chelonobacter* genus, was described in 2009 (8). Based on the phylogeny inferred from analyzing 16S rRNA, *rpoB*, *infB*, and *recN* gene sequences, *C. oris* clusters together with [*Pasteurella*] *testudinis*, a species also found in captive tortoises (9). The two taxa form a distinct cluster distantly related to all other species within the *Pasteurellaceae* family (3).

The genomic DNA from *C. oris* strain 1662^T was isolated using a blood and tissue kit (catalog no. 69506; Qiagen). The draft genome sequencing was performed at the National High-Throughput DNA Sequencing Center of the University of Copenhagen. The MiSeq instrument (Illumina) was used for the genome sequencing with a 150-bp paired-end-read format. The CLC Genomic Workbench version 6.5.1 software package (CLC, Denmark) was used to perform quality trimming and *de novo* assembly of the reads. The run yielded 1,156,558 high-quality filtered sequences in pairs containing 134,646,107 bases, which provided an average of 50-fold coverage of the genome. The assembly resulted in 35 contigs ranging from 1,370 bp to 447,435 bp in size. Prediction of protein-coding sequences and annotation was performed by the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (<http://www.ncbi.nlm.nih.gov/genomes/static/Pipeline.html>). The total estimated size of the genome was 2,610,727 bp with a G+C content of 47.1%. The genome contained 2,284 coding DNA sequences (CDSs), 3 rRNAs, and 48 tRNA sequences. In comparison to other *Pasteurellaceae* members, *C. oris* has one of the largest genomes reported and a relatively high G+C content (10).

All *C. oris* strains are β -hemolytic on calf-blood agar. To identify the potential hemolysins, we compared the assembled genome to a local database built from RTX-like proteins present in different gammaproteobacteria (11), and to the MViDB database (12)

with BLASTp. RTX toxins are the most abundant toxins among *Pasteurellaceae* bacteria, where they often act as the primary virulence factor (7). Although several proteins (locus tags: OA57_07455, OA57_11040, OA57_08905, and OA57_11380) harboring RTX repeats were identified in *C. oris*, none of these showed significant sequence similarity to known RTX toxins. Interestingly, another possible hemolysin (locus tag OA57_07415), together with a channel-forming transporter protein (locus tag OA57_07410) was identified. Both proteins showed significant sequence similarity to the Hpma (accession no. P16466.1) and Hpmb (accession no. P16465.1) proteins, respectively, of *Proteus mirabilis* (13). The Hpma protein belongs to the family of the pore-forming calcium-independent cytolysins described previously (14). Homologs of these proteins were not detected in other members of *Pasteurellaceae*, and thus their role in the pathogenicity of *C. oris* remains to be investigated.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number **JSUM00000000**. The version described in this paper is version JSUM01000000.

ACKNOWLEDGMENT

This work was supported by grants from the Faculty of Health and Medical Sciences, University of Copenhagen.

REFERENCES

1. Janda M. 2011. Update on family *Pasteurellaceae* and the status of genus *Pasteurella* and genus *Actinobacillus*. *Clin. Microbiol. Newsl.* 33:135–144. <http://dx.doi.org/10.1016/j.clinmicnews.2011.08.003>.
2. Kuhnert P, Scholten E, Haefner S, Mayor D, Frey J. 2010. *Basfia succiniciproducens* gen. nov., sp. nov., a new member of the family *Pasteurellaceae* isolated from bovine rumen. *Int. J. Syst. Evol. Microbiol.* 60: 44–50. <http://dx.doi.org/10.1099/ijs.0.011809-0>.
3. Foster G, Higgins R, Leclair D, Korczak BM, Mikaelian I, Patterson IA, Kuhnert P. 2011. Proposal of *Bisgaardia hudsonensis* gen. nov., sp. nov. and an additional genomospecies, isolated from seals, as new members of the family *Pasteurellaceae*. *Int. J. Syst. Evol. Microbiol.* 61:3016–3022. <http://dx.doi.org/10.1099/ijs.0.028027-0>.
4. Hansen MJ, Bertelsen MF, Christensen H, Bojesen AM, Bisgaard M. 2012. *Otariodibacter oris* gen. nov., sp. nov., a member of the family *Pas-*

- teurellaceae* isolated from the oral cavity of pinnipeds. *Int. J. Syst. Evol. Microbiol.* 62:2572–2578. <http://dx.doi.org/10.1099/ijs.0.039586-0>.
5. Korczak BM, Bisgaard M, Christensen H, Kuhnert P. 2014. *Frederiksenia canicola* gen. nov., sp. nov. isolated from dogs and human dog-bite wounds. *Antonie Van Leeuwenhoek* 105:731–741. <http://dx.doi.org/10.1007/s10482-014-0129-0>.
 6. Hansen M, Bertelsen MF, Christensen H, Bojesen AM. 2014. *Bisgaardia miroungae*, sp. nov., a new member of the family *Pasteurellaceae* isolated from the oral cavity of northern elephant seals (*Mirounga angustirostris*) and emended description of *Bisgaardia*. *Int. J. Syst. Evol. Microbiol.* [Epub ahead of print]. <http://dx.doi.org/10.1099/ijs.0.065060-0>.
 7. Frey J. 2011. The role of RTX toxins in host specificity of animal pathogenic *Pasteurellaceae*. *Vet. Microbiol.* 153:51–58. <http://dx.doi.org/10.1016/j.vetmic.2011.05.018>.
 8. Gregersen RH, Neubauer C, Christensen H, Bojesen AM, Hess M, Bisgaard M. 2009. Comparative studies on [*Pasteurella*] *testudinis* and [*P.*] *testudinis*-like bacteria and proposal of *Chelonobacter oris* gen. nov., sp. nov. as a new member of the family *Pasteurellaceae*. *Int. J. Syst. Evol. Microbiol.* 59:1583–1588. <http://dx.doi.org/10.1099/ijs.0.007203-0>. PubMed.
 9. Snipes KP, Kasten RW, Calagoan JM, Boothby JT. 1995. Molecular characterization of *Pasteurella testudinis* isolated from desert tortoises (*Gopherus agassizii*) with and without upper respiratory tract disease. *J. Wildl. Dis.* 31:22–29. <http://dx.doi.org/10.7589/0090-3558-31.1.22>.
 10. Kuhnert P, Korczak BM. 2006. Prediction of whole-genome DNA-DNA similarity, determination of G+C content and phylogenetic analysis within the family *Pasteurellaceae* by multilocus sequence analysis (MLSA). *Microbiology* 152:2537–2548. <http://dx.doi.org/10.1099/mic.0.28991-0>.
 11. Linhartová I, Bumba L, Mašín J, Basler M, Osička R, Kamanová J, Procházková K, Adkins I, Hejnová-Holubová J, Sadílková L, Morová J, Sebo P. 2010. RTX proteins: a highly diverse family secreted by a common mechanism. *FEMS Microbiol. Rev.* 34:1076–1112. <http://dx.doi.org/10.1111/j.1574-6976.2010.00231.x>.
 12. Zhou CE, Smith J, Lam M, Zemla A, Dyer MD, Slezak T. 2007. MvirDB—a microbial database of protein toxins, virulence factors and antibiotic resistance genes for bio-defence applications. *Nucleic Acids Res.* 35:D391–D394. <http://dx.doi.org/10.1093/nar/gkl791>.
 13. Uphoff TS, Welch RA. 1990. Nucleotide sequencing of the *Proteus mirabilis* calcium-independent hemolysin genes (*hpmA* and *hpmB*) reveals sequence similarity with the *Serratia marcescens* hemolysin genes (*shlA* and *shlB*). *J. Bacteriol.* 172:1206–1216.
 14. Welch RA. 1991. Pore-forming cytolysins of gram-negative bacteria. *Mol. Microbiol.* 5:521–528. <http://dx.doi.org/10.1111/j.1365-2958.1991.tb00723.x>.